




# ACTA

## MINERALOGICA-PETROGRAPHICA

### FIELD GUIDE SERIES

**Volume 2**

**Szeged, 2010**



BERNHARD FÜGENSCHUH & PETER TROPPER

**Transect through the Eastern Alps:  
Petrology and geology in the  
surroundings of the Brenner base tunnel**

**IMA2010 FIELD TRIP GUIDE AT2**





# **ACTA MINERALOGICA-PETROGRAPHICA**

*established in 1923*

## **FIELD GUIDE SERIES**

HU ISSN 0324-6523

HU ISSN 2061-9766

### **Editor-In-Chief**

*Elemér Pál-Molnár*

*University of Szeged, Szeged, Hungary*

*E-mail: palm@geo.u-szeged.hu*

### **EDITORIAL BOARD**

Péter Árkai, György Buda, István Dódon, Tamás Fancsik, János Földessy, Szabolcs Harangi, Magdolna Hetényi, Balázs Koroknai, Tivadar M. Tóth, Gábor Papp, Mihály Pósfai, Péter Rózsa, Péter Sipos, Csaba Szabó, Sándor Szakáll, Tibor Szederkényi, István Viczián, Tibor Zelenka

### **Guest Editor of this Volume**

Gábor Papp

*Hungarian Natural History Museum, Budapest, Hungary*

*E-mail: pappmin@ludens.elte.hu*



This volume was published for the  
375<sup>th</sup> anniversary of the  
Eötvös Loránd University, Budapest.



The publication was co-sponsored by the  
Eötvös University Press Ltd., Budapest.

IMA2010 ([www.ima2010.hu](http://www.ima2010.hu)) is organised in the frame of the ELTE375 scientific celebration activities.

### **IMA2010 FIELD TRIP SUBCOMMITTEE**

**Chairmen:** Friedrich Koller, University of Vienna (AT) and Ferenc Molnár, Eötvös L. University, Budapest (HU)

**Members:** Volker Höck, University of Salzburg (AT); Corina Ionescu, Babeş-Bolyai University, Cluj-Napoca (RO); Veselin Kovachev, Sofia University "St. Kliment Ohridski" (BG); Marek Michalik, Jagellonian University, Kraków (PL); Milan Novák, Masaryk University, Brno (CZ); Ladislav Palinkaš, University of Zagreb (HR); Simona Skobe, University of Ljubljana (SI); Sándor Szakáll, University of Miskolc (HU); Pavel Uher, Comenius University, Bratislava (SK); Nada Vasković, University of Belgrade (RS)

### **OFFICERS OF THE IMA2010 ORGANISING COMMITTEE**

**Chairman:** Tamás G. Weiszbürg, Budapest, Hungary,

**Secretary General:** Dana Pop, Cluj-Napoca, Romania

### **Editorial Office Manager**

Anikó Batki

*University of Szeged, Szeged, Hungary*

*E-mail: batki@geo.u-szeged.hu*

### **Editorial Address**

*H-6701 Szeged, Hungary*

*P.O. Box 651*

*E-mail: asviroda@geo.u-szeged.hu*

The Acta Mineralogica-Petrographica is published by the Department of Mineralogy, Geochemistry and Petrology, University of Szeged, Szeged, Hungary

© Department of Mineralogy, Geochemistry and Petrology, University of Szeged

ISBN 978-963-306-034-6


---

*On the cover: View towards North from Brenner pass area showing the Brenner fault (valley) together with the Tauern window footwall (right side) and Ötztal hangingwall (left side). Northern Calcareous Alps in the background.*



# Geological cross section along the planned Brenner Base Tunnel

Töchterle, A., Brandner, R., Pomella, H. & Reiter, F.



**Tectonic units and marker units**

Quaternary	Triassic at the base of the Lower Bündnerschiefer	Furtschagl schists
Patscherkofel Basement Unit	Kaserer Fm.	Tulfer-Senges Unit and Greiner Series
Innsbruck Quartzphyllite	Hochstegen Fm. (Upper Jurassic), kyanite-quartzite (?Middle Jurassic)	"Maulser Trias" (Austroalpine Permo-Mesoz. U.)
Innsbruck Quartzphyllite, higher metamorphic	Aigerbach Fm. (Triassic)	Austroalpine basement south of the Tauern Window (incl. quest. U. Carboniferous)
Tarntal Permo-Mesozoic Units	Wustkogel Fm. (Permo-Triassic)	Oligocene calcalkaline intrusives
Bündnerschiefer undifferentiated	Central Gneiss and "Altes Dach" (Variscan basement)	Brixen Granite
Upper Bündnerschiefer ("Tauernflysch")		
Lower Bündnerschiefer		

**Structures**

brittle fault	reverse fault
ductile fault	strike-slip (oblique) fault
fault inferred	detachment
remote sensing	thrust fault
major fault	axial plane - gently and tight fold
minor fault	axial plane - isoclinal fold

**Mass movements**

deep-seated gravitational slope deformation

**Drillings (partly projected)**

Drillings (partly projected)

Fig. 1. Transect through the Eastern Alps showing the Brenner base tunnel traverse (from Brandner *et al.*, 2008)

The shallow part of the section is based on the partly revised longitudinal section 1:10.000 for the Brenner Base Tunnel (Phase II). This was a consortium project of the Austrian Geological Survey, the Institute for Geology and Paleontology from the University of Innsbruck and the "Consorzio Ferrara Ricerche". We are very grateful to the Brenner Base Tunnel Company (BBT SE) for funding this project.





# Geological cross section along the planned Brenner Base Tunnel

Author: G. Zischler, M. G. Zischler, F. Zischler



Geological cross section along  
the planned Brenner Base Tunnel





X 175526





# **Transect through the Eastern Alps: Petrology and geology in the surroundings of the Brenner base tunnel**

**BERNHARD FÜGENSCHUH<sup>1</sup>\* AND PETER TROPPER<sup>2</sup>**

<sup>1</sup> Bernhard Fügenschuh, Institut für Geologie und Paläontologie, Universität Innsbruck, Innrain 52, A-6020 Innsbruck, Austria; bernhard.fuegenschuh@uibk.ac.at, \*communicating author

<sup>2</sup> Peter Tropper, Institut für Mineralogie und Petrographie, Universität Innsbruck, Innrain 52, A-6020 Innsbruck, Austria; peter.tropper@uibk.ac.at

## **Table of contents**

1. Introduction .....	2
2. Tectonic and metamorphic evolution .....	2
3. Tectonic overview .....	2
4. Field stops .....	6
Day one .....	6
4.1 Field stop 1: Entrance of the Brenner base tunnel in Innsbruck, Sillschlucht .....	6
4.2 Field stop 2: Cataclasites of Brenner fault zone, Stefansbrücke .....	6
4.3 Field stop 3: Overview of regional geology, Nösslach .....	7
4.4 Field stop 4: Brenner fault mylonites, Zentralgneiss, Brenner Pass .....	7
4.5 Field stop 5: Periadriatic line, Oligocene intrusive body, Mauls .....	7
4.6 Field stop 6: Permian intrusive body, Mittewald .....	8
Day two .....	8
4.7 Field stop 7: Jaufen fault, Cretaceous tectonics, Jaufen Pass .....	8
4.8 Field stop 8: Southern Ötztal Complex, Cretaceous eclogite-facies, Saltaus .....	8
4.9 Field stop 9: Ötztal/Schneeberg Complex, Cretaceous metamorphism, Timmelsjoch .....	8
4.10 Field stop 10: Eclogites near Längenfeld, Variscan eclogite facies, Burgstein .....	9
4.11 Field stop 11: Köfels rock fall, Köfels .....	9
5. References .....	9
Appendix – Itinerary for IMA2010 AT2 Field trip .....	13

## **1. Introduction**

The visited area highlights the polyphase tectonic and metamorphic evolution of the eastern Alps along the Brenner base tunnel transect (Fig. 1). While the Austroalpine units bear substantial information related to the Cretaceous evolution/orogeny, the Tauern window exposes units and rocks of European

provenance, metamorphosed and deformed during Tertiary collision. With additional stops in the Southern Alps the excursion provides insight in the pre-Alpine evolution of this area. During the Neogene the final shaping of this area occurred in the context of lateral extrusion and exhumation.



## 2. Tectonic and metamorphic evolution

The Permian evolution of the eastern Alps is highly influenced by magmatic activity represented by both, intrusive and extrusive rocks. This activity plays a key role in the Mesozoic evolution of this area, characterized by the opening of Neotethys in the east and a continuously subsiding passive margin, surrounding the Neotethys embayment. While in mid-Jurassic times the Alpine Tethys opened in the west (central Atlantic rifting), the Neotethys ocean underwent a complex evolution of subduction/obduction and re-rifting, starting in the late Jurassic (Fig. 2). It is this late Jurassic/early Cretaceous subduction in the context of which the Austroalpine units underwent metamorphic overprint up to eclogite-facies conditions. While the proper suture of this event is currently not exposed (most likely to be found under the Neogene Pannonian basin), the eclogite corridor of the eastern Alps marks the trace of the intrabasement collision between the two opposing sides of the former embayment. Cretaceous collision and metamorphism in the eastern Alps was followed by late Cretaceous extension and exhumation of high-grade rocks. This phase is well known in the Alps but also in the Carpathians and Dinarides as the Gosau phase, which marks the end of the Cretaceous cycle in the eastern Alps. The further evolution is triggered by subduction of the Alpine Tethys and final collision of the European and Adriatic plate in the Tertiary. In the course of the Tertiary orogeny the European lower plate was partly subjected to eclogite-facies metamorphism.

In the Eastern Alps, the Austroalpine basement typically shows polymetamorphism due to a sequence of metamorphic overprints, affecting this part of the Alps (Oberhänsli *et al.*, 2004; Schuster *et al.*, 2004). The most dominant metamorphic overprints are the Variscan- and the Eo-Alpine metamorphic event (Hoinkes *et al.*, 1999; Neubauer *et al.*, 1999; Thöni, 1999). In recent years, geochronological data also point to a widespread Permian

thermal overprint mainly observed in the eastern part of the Austroalpine units (Schuster *et al.*, 2001). Since Paleozoic times it was affected by four regional metamorphic events (Oberhänsli *et al.*, 2004; Schuster *et al.*, 2004) in Late Devonian to Carboniferous, Permian, Cretaceous and Oligocene to Miocene times. The Late Devonian to Carboniferous imprint is related to the Variscan orogenic cycle. In Permian times large parts of the Austroalpine units were affected by lithospheric extension and related high-temperature/low-pressure metamorphism. The Eo-Alpine tectonothermal event in the Cretaceous is due to intracontinental shortening within the Austroalpine unit. Finally in Oligocene to Miocene times a thermal influence of the Alpine metamorphic event, related to the continental collision after the closure of the Penninic oceans can be recognised in the tectonically lowermost Austroalpine units to the south of the Tauern Window.

Based on their position within the Alpine orogen the individual lithostratigraphic units of the Austroalpine unit experienced different metamorphic histories during these four metamorphic events mentioned above. Most of the units are

polymetamorphic, but several of them experienced only one imprint: e.g. in the Silvretta-Seckau-, Ötztal-Bundschuh- and Drauzug-Gurktal nappe systems, as well as in the Greywacke zone units with a dominating Variscan metamorphic imprint occur. In many cases they are transgressively overlain by unmetamorphosed or very-low grade metamorphic Permian-Mesozoic sequences, whereas others experienced an additional Eo-Alpine overprint. Most of the units in the Koralpe-Wölz nappe system experienced a Permian-Triassic imprint and an Eo-Alpine overprint, which reached up to eclogite facies conditions. However, parts of the Wölz- and Radenthein Complexes show only the Eo-Alpine metamorphic imprint.

## 3. Tectonic overview

In the following a brief description of the different units as shown in Figure 2 will be given:

The **Tauern Window** (Fig. 3, units 32–35) exposes Penninic and sub-Penninic (European distal margin) units in the footwall of the Austroalpine nappe complex, which forms the hanging-wall plate during




Fig. 2. Paleogeographical map displaying the relevant oceans for the Alps (after Schmid *et al.*, 2004).






Fig. 3. Tectonic map of the Brenner area (from Schmid *et al.*, 2004): 12: upper crust Southern Alps; 13: post-Variscan cover Southern Alps; 16, 17, 18: Upper Austroalpine nappes (NCA and GWZ); 21: Upper Austroalpine basement nappe ÖC; 22: Upper Austroalpine basement nappe Texel complex; 23: Upper Austroalpine basement nappe (Campo, Innsbruck quartzphyllite); 24: Lower Austroalpine basement nappe; 26: Upper Penninic nappe; 32: Lower Penninic nappe; 33: Sub-Penninic: Mesozoic cover; 34: Sub-Penninic: non eclogitic basement; 35: Sub-Penninic: eclogitic basement.

Tertiary plate collision. The units within the Tauern Window are separated into several nappes that are characterized by typical lithofacial assemblages. From the footwall to the hanging-wall, the nappe stack includes:

- (1) The Venediger Nappe and the Wolfendom Nappe comprising a pre-Variscan basement intruded by Variscan granitoids (Zentralgneis cores) and a cover sequence of Jurassic metacarbonates, and Cretaceous metapelites and metapsammities.
- (2) The Storz and Riffel Nappes comprising Variscan and Alpidic polymetamorphic basement rocks covered by the late Palaeozoic(?) or Cretaceous(?) metapelites and graphitic quartzites.
- (3) The Eclogite Zone which is restricted to the central southern Tauern Window and which is characterized by a Mesozoic volcano-sedimentary sequence.
- (4) The Rote Wand-Modereck Nappe, consisting of basement rocks which are covered by Permian to Triassic quartzites and Triassic metacarbonates, Jurassic breccias, calcareous micaschists and metatuffs as well as Cretaceous metapelites and metapsammities.
- (5) The Glockner Nappe, which contains an oceanic basement (serpentinites and ultramafic rocks) and a partly incomplete ophiolitic sequence.
- (6) The Matrei Zone, interpreted to represent an accretionary wedge, is characterized by metamorphic flysch sediments, breccias and olistolithes mainly of Austroalpine derivation.

- (7) The Klammkalk Zone, consisting of calcareous schists, massive marbles and minor phyllites, forming a low-grade metamorphic equivalent to the 'Bündnerschiefer' in the Glockner Nappe.

These units are surrounded by the remnants of Lower Austroalpine units. The Eclogite Zone contains mafic eclogites of tholeiitic and slightly alkaline chemical composition (Miller, 1974, 1977, 1987). The protoliths are assumed to be basalts with an intra-plate character (Höck & Miller, 1987). They are often retrogressed to garnet-amphibolites and garnet-bearing greenschists due to a Barrovian-type, greenschist to amphibolite-facies overprint. The associated metasediments experienced the same high-pressure metamorphism (Franz & Spear, 1983; Dachs, 1986, 1990; Spear & Franz, 1986). The rocks exposed within the Eclogite Zone experienced a polyphase metamorphic evolution. Starting with inclusions in garnets, which are sometimes interpreted as pseudomorphs after lawsonite, a first stage of metamorphism at ca. 400 °C can be deduced (Miller, 1977, 1986; Frank *et al.*, 1981, 1987). Eclogite-facies metamorphism is only rarely documented and is only observed clearly in the Eclogite Zone. The eclogite-facies rocks were buried to a depth of at least 65 km (20 kbar, 600 °C; Holland, 1979; Dachs, 1986, 1990; Frank *et al.*, 1987a; Droop *et al.*, 1990; Selverstone *et al.*, 1992; Zimmermann *et al.*, 1994; Getty & Selverstone, 1994). On the other hand, the entire nappe pile of the Tauern Window was subsequently affected by pervasive blueschist-facies metamorphism with *P-T* conditions ranging from 7–15 kbar and ca. 450 °C (Raith *et al.*, 1980; Holland, 1979; Zimmermann *et al.*, 1994; Holland & Richardson, 1979; Selverstone *et al.*, 1992; Cliff *et al.*, 1985; Droop, 1985; Holland & Ray, 1985; Frank *et al.*, 1987; Selverstone, 1993). Finally, the entire nappe pile was affected by Barrovian-type upper greenschist to lower amphibolite-facies metamorphism (e.g., Frank *et al.*, 1987a; Selverstone, 1993). Phengite <sup>40</sup>Ar/<sup>39</sup>Ar mineral ages of ca. 36–32 Ma (Zimmermann *et al.*, 1994) from the Eclogite Zone are interpreted to represent cooling ages subsequent to Eocene blueschist-facies metamorphism (Zimmermann *et al.*, 1994).

The **Innsbruck Quartzphyllite** (Fig. 3, unit 23) crops out between Mittersill in the east and Innsbruck in the west. It is typically a rather monotonous, fine-grained, greenish to grayish phyllitic schist, with the mineral assemblage muscovite + chlorite + albite + quartz ± calcite. Locally, garnet-bearing schists occur south of the Patscherkofel. It has been divided into three stratigraphical units consisting of Devonian carbonatic black shales, Silurian carbonatic-sericitic phyllites and Ordovician quartzphyllites and greenschists, however numerous transitions may be found (Haditsch & Mostler, 1982). Although most of the Innsbruck Quartzphyllites were affected by lower greenschist-facies metamorphism (Hoschek *et al.* 1980; Sassi & Spiess, 1992; Piber & Tropper, 2005), some central parts of the Innsbruck Quartzphyllite have been affected by middle greenschist-facies metamorphism (Kolenprat *et*



al. 1999; Piber & Tropper, 2007). Geochronological investigations revealed a complex metamorphic history indicating a possible Permian- and Eo-Alpine overprint (Dingeldey *et al.*, 1997; Rockenschaub *et al.*, 1999; Handler *et al.*, 2000). Recently, a number of new results have been obtained concerning the internal structure of the Innsbruck Quartzphyllite (Kolenprat *et al.*, 1999). Large parts of the Innsbruck Quartzphyllite must therefore be considered as highly deformed, retrograde old (Variscan?) basement. These studies revealed a metamorphic zonation with garnet-free phyllites at the northern and southern rims and garnet bearing phyllites in the central part, thus reflecting a slightly higher grade of metamorphism in the center. This observation was interpreted in terms of a km-scale isoclinal fold of the Innsbruck Quartzphyllite (Schmidegg 1964, Rockenschaub 1998; Kolenprat 1998). Kolenprat *et al.* (1999) show, that the Innsbruck Quartzphyllite has a complex deformation history, with structures ranging from pre-Alpine (Variscan) to late Alpine (Neogene) in age. The pre-Alpine foliation is preserved only locally. During the Eo-Alpine orogeny, intensive mylonitization associated with W- to NW-directed nappe stacking, occurred. The Meso- and Neo-Alpine deformation is characterized by the imbrication of Austroalpine units as a consequence of N-directed thrusting of the Austroalpine nappes over the Penninic Units and subsequent exhumation of the Tauern-Window during N-S-shortening and E-W-extension (Kolenprat *et al.*, 1999). On top of the Innsbruck Quartzphyllite, two units occur in the vicinity of the Brenner base tunnel, namely the Patscherkofel Crystalline Complex and the Kellerjochgneiss.

The **Patscherkofel Crystalline Complex** (PCC) is part of the Austroalpine basement nappes north of the Tauern Window, which is tectonically located on top of the Innsbruck Quartzphyllite. The PCC is mainly composed of mica schists with the mineral assemblage plagioclase + muscovite + biotite + chlorite + quartz  $\pm$  chloritoid  $\pm$  garnet<sub>1,2</sub>  $\pm$  ilmenite  $\pm$  clinozoisite  $\pm$  staurolite  $\pm$  margarite. Garnet<sub>1</sub> + staurolite represent Pre-Alpine relicts, all other minerals are part of the Eo-Alpine mineral assemblage. Thermobarometric investigations of Piber *et al.* (2008) yielded temperatures between 510 °C and 570 °C and pressures ranging from 9.5 to 12.2 kbar for the samples from the PCC.

The **Kellerjochgneiss** or Schwaz Augengneiss is also part of the Austroalpine basement nappes north of the Tauern Window. The Kellerjochgneiss is a former I-type augengneiss and contains the mineral assemblage muscovite + plagioclase + chlorite + quartz  $\pm$  biotite  $\pm$  clinozoisite. It tectonically overlies the Innsbruck Quartzphyllite. Multiequilibrium calculations of samples of the Kellerjochgneiss yielded pressures ranging from 3.2 to 6.8 kbar and temperatures ranging from 285 to 345 °C (Piber, 2005).

The **Tarntal Nappe** (Fig. 3, unit 24) is a complex geologic unit, which consists of low-grade metamorphic sediments and

a dismembered ophiolite body, and is tectonically emplaced between the contact of the Austroalpine Innsbrucker Quartzphyllite nappe and the Penninic Glockner Nappe at the northern margin of the Tauern Window (Tirol, Austria). During the Alpine metamorphic overprint, blueschists formed at the contact between the metasedimentary units and the ultramafic units of the ophiolite within the Tarntal Nappe. Dingeldey *et al.* (1997) and Klier (2005) obtained metamorphic peak *P-T* conditions of 350 °C and 8 to 10.5 kbar. The age of the metamorphic overprint was dated with 50–40 Ma depending on the tectonic position of the samples.

The Wildschönau Schists and the Schwaz Dolomite form part of the **Greywacke Zone** (unit 18). According to Mostler (1973) the Western Greywacke Zone is a stratigraphic sequence of metasediments with volcanic intercalations, ranging from the Ordovician to the Late Devonian. The Wildschönau Schists are composed of light gray phyllites similar to the Innsbruck Quartzphyllite. Roth (1983) characterized two different varieties of the Wildschönau Schists, the sandy type and the phyllitic type. The mineral assemblage of the Wildschönau Schists is very similar to the Innsbruck Quartzphyllites containing the mineral assemblage muscovite + chlorite + albite + quartz  $\pm$  calcite. In the Western Greywacke Zone, modern thermobarometric data were lacking until recently. Based on index minerals, Hoschek *et al.* (1980) estimated lower greenschist-facies conditions for the Wildschönau Schiefer. Piber (2005) obtained *P-T* conditions of 4.5 kbar and 330 °C, based on multi-equilibrium thermobarometry on one sample of the Wildschönau Schists. Geochronological investigations by Handler *et al.* (2000) and Anglmeier *et al.* (2000) indicate a Permian metamorphic overprint. Using <sup>40</sup>Ar/<sup>39</sup>Ar and <sup>87</sup>Rb/<sup>87</sup>Sr recently yielded Eo-Alpine ages in the central Greywacke Zone of 102–98 Ma (Schmidlechner *et al.*, 2006) in addition to the <sup>87</sup>Rb/<sup>87</sup>Sr ages of 137 to 127 Ma and <sup>40</sup>K/<sup>39</sup>Ar ages of 113 to 92 and 113 to 106 Ma from the Greywacke Zone close to Zell am See which also give reasonable evidence for an Eo-Alpine metamorphic overprint around ca. 300 °C (Kralik *et al.*, 1987).

The **Ötztal Complex** (Fig. 3, ÖC, unit 21) is a large crystalline complex in the western part of the Austroalpine units. The ÖC consists of quartzofeldspathic and metapelitic metasediments with various intercalations of orthogneisses, amphibolites and rare metacarbonates. The oldest metamorphic event is “Caledonian” in age (460–490 Ma), leading to the formation of orthogneisses (Thöni, 1986) and scattered occurrences of migmatites (Söllner *et al.*, 1982; Söllner & Schmidt, 1981; Klötzli-Chowanetz *et al.*, 1997). Hoinkes (1973) estimated the *P-T* conditions of migmatite formation with 680 °C and  $\geq 4$  kbar, Thöni *et al.* (2008) obtained slightly lower pressures of  $< 2.8$  kbar. The Variscan metamorphic overprint ranges from 390–295 Ma (Thöni, 1999). The first stage of the Variscan event was a high-pressure metamorphism around 373–359 Ma, leading to the formation of eclogites in the central part of



the ÖC (Miller & Thöni, 1995). The conditions of the eclogite facies were estimated to be 710–750 °C and 27–28 kbar (Miller & Thöni, 1995). The dominant amphibolite facies metamorphism occurred around 330–350 Ma, as evident from Sm–Nd garnet-whole rock ages from micaschists (Schweigl, 1993, 1995; Hoinkes *et al.*, 1997; Thöni, 1999). Purtscheller (1969) observed on the basis of the regional distribution of  $\text{Al}_2\text{SiO}_5$  polymorphs in metapelites a systematic regional zonation and distinguished within the ÖC two zones: (1) the southern and northern kyanite zone and (2) the central sillimanite zone including the andalusite zone in the west. Hoinkes & Thöni (1993) pointed out that the occurring mineral zonation does not have to be the consequence of a single Variscan metamorphic event. Tropper & Hoinkes (1996) estimated  $P$ – $T$  conditions of 570–640 °C and 5.8–7.5 kbar for the northwestern part of the ÖC. The youngest metamorphic event in this Austroalpine basement occurred during the Cretaceous Eo-Alpine orogeny (100–73 Ma, Thöni, 1981; Thöni, 1999; Exner *et al.*, 2001; Habler *et al.*, 2001). The intensity of the Eo-Alpine overprint varies within the ÖC and increases from NW (lower greenschist facies) to SE (epidote-amphibolite facies) and reaches 550–600 °C and  $\geq 11$  kbar in the Schneeberg Complex (Hoinkes *et al.*, 1991; Konzett & Hoinkes, 1996). This zonation ends abruptly at the Passeier – Jaufen Line. This leads also to resetting of Variscan cooling ages from the NW to the SE (Thöni 1981; Thöni, 1999). To the south of the Passeier–Jaufen Line, in the Meran–Mauls basement (MMB) only a weak Eo-Alpine metamorphic overprint can be detected (Spiess, 1995).

The Paleozoic **Schneeberg Complex (SC)** consists of at least three narrow and structurally complicated synclines folded into the southern part of the polymetamorphic ÖC. The metasediments of the SC are characterized by the frequent presence of metamarls and marbles as opposed to the ÖC where such lithologies are missing with few exceptions known from the central ÖC (Hoinkes & Thöni, 1993). Sölvä *et al.* (2001, 2005) define the SC as a tectonic unit associated with the Schneeberg normal fault zone, separated from the ÖC and strongly affected by the Eo-Alpine tectonometamorphic event. Most recently, the discovery of Eo-Alpine andalusite coexisting with staurolite + biotite + garnet at temperatures  $\geq 540$  °C places tight constraints on the retrograde part of the Eo-Alpine  $P$ – $T$  path of SC rocks that is characterized by an almost isothermal decompression from the peak of metamorphism into the andalusite stability field. SC rocks do not provide any evidence for an earlier, high-pressure stage. Geochronologic results by Elias (1998) and Fügenschuh *et al.* (2000) point to a rapid uplift and exhumation of both SC and underlying ÖC after the peak of metamorphism. According to Fügenschuh *et al.* (2000) initial uplift rates were around 1 mm/year starting around 90 Ma, which decreased to values of 0.2 to 0.7 mm/year (Hoinkes *et al.*, 1991).

The Variscan basement of the **Southalpine domain** (Fig. 3, unit 13) is confined to the west and north by the Periadriatic (Giudicarie line and Pustertal line) fault system. In the south-east, small basement outcrops within the Cenozoic molasse deposits of the Po Plain occur. Most of the basement is comprised of monotonous quartzphyllites (Brixen Quartzphyllite), which were pervasively affected by the Variscan metamorphic and tectonic overprint. Due to its now slightly tilted position, the basement shows a metamorphic gradient, which increases from southeast towards northwest (Sassi & Spiess, 1993). In the area of Toblach, the basement contains the mineral assemblage quartz + chlorite + white mica + albite and represents the lowest peak metamorphic conditions of the basement with temperatures of 350–400 °C and a pressure of ca. 0.4 GPa. The metamorphic conditions increase towards the northwest and reach maximum  $P$ – $T$  conditions in the area of Brixen/Bressanone. In this area the basement contains the mineral assemblage quartz + biotite + chlorite + white mica + garnet + albite + plagioclase and calculated  $P$ – $T$  conditions based on garnet-biotite thermometry and plagioclase-biotite-garnet-muscovite barometry yielded temperatures of 450–550 °C and pressures of 0.5–0.65 GPa (Ring & Richter 1994).

In the Southalpine domain the **Permian intrusive complexes** of the Brixen granite, Ifinger granodiorite and Kreuzberg granite cover an area of  $\sim 250$  km<sup>2</sup> and are thought to have been the result of the collapsing Variscan orogenic belt, which led to the formation of large extensional terrains (Acquafredda *et al.*, 1997; Bargossi *et al.*, 1981; Bargossi *et al.*, 1998; Del Moro & Visona, 1982). Although these intrusive complexes were already mapped (Del Moro & Visona, 1982) and considerable literature concerning their magmatic evolution already exists (Bonin *et al.*, 1993; Schuster *et al.*, 2001; Dal Piaz & Martin, 1998) almost no petrological (Bonin *et al.*, 1993; Acquafredda *et al.*, 1997) and only few geochronological data (Bonin *et al.*, 1993; Borsi *et al.*, 1972; Rottura *et al.*, 1998) are available so far. The investigated intrusions namely the Brixen/Bressanone granodiorite, the Ifinger/Ivigna granite and the Kreuzberg/Monte Croce granite are all aligned along the Periadriatic Lineament. The major part of the Brixen granite is granitic to granodioritic in composition with the mineral assemblage K-feldspar + plagioclase + biotite + quartz + accessories (zircon, apatite, ilmenite  $\pm$  monazite). Zircon ages from three intrusions in the Southalpine basement show ages ranging from  $293 \pm 12$  Ma (Kreuzberg granite),  $289 \pm 6.1$  Ma (Ifinger granodiorite) to  $278 \pm 12$  Ma (Brixen granite). These geochronological data seem to indicate a rejuvenation trend from south-west to north-east which might be evidence for a plane motion of the north-western realm of the Athesian Volcanic Group (AVG) caldera during the late Variscan orogenic collapse (Thöni, 2008).



## 4. Field stops

### Day one

#### 4.1 Field stop 1: Entrance of the Brenner base tunnel in Innsbruck, Sillschlucht

Located at the southern limit of Innsbruck directly under the ski jump stadium this stop allows to discuss the local geology in the area of the northern entrance of the Brenner base tunnel (Fig. 4). The outcrops are located along a small track near the river Sill. Apart from unstable Quaternary deposits, giving rise to mobility right at the tunnel entrance, the local geology is dominated by quartzphyllite. Metamorphosed under lower greenschist-facies conditions during the Variscan cycle the rocks show multiphase tectonic overprint from ductile deformation during pre-Alpine and Alpine times to Neogene brittle deformation. Second phase folding led to the formation of a large recumbent fold, which has granitic basement in its core. Although rather monotonous, different lithologies can be observed along the track, ranging from porphyries to amphibolites/prasinites to marbles. According to the overall appearance the quartzphyllite unit most likely represents a metamorphosed turbidite sequence.

#### 4.2 Field stop 2: Cataclasites of Brenner fault zone, Stefansbrücke

Some 5 km south of Innsbruck along the federal road to Brennerpass brittle fault rocks related to the Brenner Fault Zone are outcropping in an abandoned quarry. The tectonically overprinted host rock is the basement of the ÖC, various stages of brittle deformation can be studied. While the least deformed part still displays the original features (schistosity, folds in paragneiss) deformation intensity increases to finally expose a fault gouge. Depending on the outcrop conditions (due to the brittle deformation they are very erodable) shear sense criteria pointing to top to the west kinematics can be observed (Fig. 5). Several similar outcrops of fault gouge in the wider Brenner area have been sampled by Zwingmann & Mancktelow (2004) for conventional K/Ar dating of clay minerals. While most of the samples yielded Neogene ages, in accordance with the brittle faulting along the BFZ, the samples from this outcrop are dominated by Cretaceous ages. This clearly points to inherited information from the overwhelming Cretaceous metamorphic overprint within the ÖC.




Fig. 4. Detailed geological map of the northern entrance of the Brenner base tunnel (Brandner *et al.*, 2008). Hatched area: rock slide; brown: quartzphyllite; red: porphyry; green: prasinite; blue: marbles.






Fig. 5. Simplified block diagram illustrating the Brenner fault zone and the involved units (Fügenschuh *et al.*, 1997).

#### 4.3 Field stop 3: Overview of regional geology, Nösslach

This overview stop is located south of the village of Steinach along a side road close to the highway exit Nösslach. If the weather is fine this stop offers a beautiful overview into the local nappe pile, exposing, from bottom to top, Zentralgneiss, Schieferhülle, Tarntaler Mesozoic, Innsbrucker Quartzphyllite, Northern Calcareous Alps, Ötztal Stubai Complex and Steinacher nappe. It is also from this point that you can judge the substantial amount of tectonically omitted material due to normal faulting along the BFZ (Fig. 5).

#### 4.4 Field stop 4: Brenner fault mylonites, Zentralgneiss, Brenner Pass

From the Brenner Pass on the Austrian side a small paved road brings us up to an active quarry in the Zentralgneiss. Along the road highly mylonitized rocks of the upper Schieferhülle (Bündnerschiefer) with beautiful shear sense criteria can be observed. In the quarry the Permian Zentralgneiss is tectonically overlain by Mesozoic strata including the limestone mylonites of the Hochstegen formation. The granitic gneiss is exposed in the core of a major WSW–ENE trending antiform, related to Tertiary nappe piling and overprinted by Neogene normal faulting (Fig. 6). Deformation within the granite is varying;

ranging from almost undeformed to strongly flattened. This allows for a guess on the width of the BFZ shear zone on the order of 1–2 km. Apart from magmatically zoned feldspars various enclaves and xenoliths can be studied.

#### 4.5 Field stop 5: Periadriatic line, Oligocene intrusive body, Mauls

In the village of Mauls (south of Sterzing) we follow a paved road towards the east entering a small valley. In the riverbed of the “Nöckebech” we can touch the contact between the metamorphosed Austroalpine basement and the unmetamorphosed southern Alpine basement together with a small Oligocene tonalite body. This part of the Periadriatic fault, a first order fault system in the Alps, does not, as elsewhere, represent the suture between Europe and Adria but forms a repeatedly activated fault zone which originally might have formed the retro-wedge of the Cretaceous orogen. The outcrop is of prime importance for the construction of the Brenner base tunnel since this tectonic zone will be crossed. On the surface it has a clear brittle character yet mylonites can definitively be expected at somewhat deeper levels. Due to its rather vertical dip it is not monitored in the seismic Transalp section which gave rise to rather provocative models. This part of the Periadriatic fault system is furthermore of great importance for solving the question of an originally straight vs. an already curved fault zone.



## 4.6 Field stop 6: Permian intrusive body, Mittewald

The major part of the Permian Brixen granodiorite is granitic to granodioritic in composition with the mineral assemblage K-feldspar + plagioclase + biotite + quartz + accessories (zircon, apatite, ilmenite  $\pm$  monazite). The south-eastern border is characterized by the occurrence of tonalite, quartz-diorite and quartz-gabbro. At the north-western rim, several unusual types of plutonic bodies occur, namely a garnet-fayalite-bearing granite near Mittewald and a two mica – andalusite – cordierite granite at the Sattelspitze. The two mica – andalusite – cordierite granite from the Sattelspitze for instance represents a peraluminous pegmatite of the latest stage of magmatic activity. The outcrop next to the road shows the rocks of the Brixen granite as well as the garnet-fayalite-bearing granite. Geochronological investigations using Sm/Nd garnet-whole rock isochron ages yielded  $280 \pm 2$  Ma (Thöny, 2008).

### Day two

## 4.7 Field stop 7: Jaufen fault, Cretaceous tectonics, Jaufen Pass

From Sterzing a spectacular road climbs up towards west to the Jaufen Pass. The road roughly follows the Jaufen fault zone, separating Cretaceous metamorphic Austroalpine basement (ÖC) from Austroalpine units which lack substantial metamorphic overprint (Meran–Mauls Basement). Across the fault zone a jump in cooling ages occurs from Cretaceous in the NW to Variscan in the SE. Thus the fault has to have a Cretaceous history. Yet recent studies (Viola *et al.* 2004; Pomella *et al.*, 2008) revealed a polyphase history, the last activity of which occurred in the course of normal faulting along the BFZ. Along the road metamorphic basement of the ÖC as well as the Meran–Mauls basement can be studied together with some outcrops of Jaufen fault mylonites.

## 4.8 Field stop 8: Southern Ötztal Complex, Cretaceous eclogite-facies, Saltaus

In this locality there is an outcrop of a few m in size along a forest track that branches off the main road from St. Martin – Meran approx. 3 km to the south of St. Martin. Metabasites of the southern Ötztal Complex hitherto mapped as amphibolites, were identified as eclogites in this outcrop. Primary mineral parageneses are tschermakitic to pargasitic green amphiboles, omphacite, garnet, phengite, zoisite, rutile and quartz. Al-pargasite forms between garnet and omphacite and is interpreted as a retrograde reaction product. In addition, retrogression of the eclogitic parageneses reflecting decreasing pressure and increasing temperature conditions are the formation of sym-

plectites of diopside and plagioclase after omphacite, Al- and Na-poor green amphiboles, grossular-poor garnet and symplectites of biotite and plagioclase replacing phengite. Minimum pressure and temperature conditions of the eclogite stage are 11–12 kbar and 500–550 °C. Maximum temperature conditions of the post-eclogitic stage were between 600 °C and 650 °C. The presence of these eclogitic metabasites as lenticular interlayers within ortho- and paragneisses indicates high-pressure metamorphic conditions within the entire rock sequence. This interpretation is confirmed by the occurrence of phengite-rich micas in orthogneisses indicating pressures of approx. 11 kbar. Secondary chemical changes of these phengites to muscovite-rich compositions again show the decreasing pressure conditions in the southern Ötztal basement after the eclogite stage.

## 4.9 Field stop 9: Ötztal/Schneeberg Complex, Cretaceous metamorphism, Timmelsjoch

At an altitude of 2474 m at the Austrian–Italian border, rocks outcropping in the immediate vicinity of the customs building are gneisses of the ÖC that form the northern border of the SC. The gneisses were affected by both Variscan and Eo-Alpine metamorphism: The Variscan metamorphism reached conditions of the amphibolite facies which led to the formation of an assemblage plagioclase + biotite + muscovite + garnet + quartz  $\pm$  staurolite  $\pm$  kyanite. The subsequent Eo-Alpine event in this area reached *P–T* conditions slightly below the stability of staurolite. This caused a retrogressive breakdown of staurolite to paragonite + chlorite + quartz. This reaction caused the formation of mica + chlorite + quartz pseudomorphs after lath-shaped staurolite crystals that may reach several cm in length and can be found on cleavage planes. *P–T* conditions sufficient for Eo-Alpine staurolite formation, however, were reached within the northernmost part of the SC.

On the Italian side at an altitude of 2200 m, after the second 180° road turn, the northern border of the main syncline of the SC towards the underlying ÖC is exposed along the roadside over a distance of approx. 100 m. At this location the distinction between both units is unambiguous due to the difference in lithologies since the rocks of the ÖC are monotonous gneisses with abundant small ( $\leq 1$  mm) garnet and biotite and the rocks of the SC are garnet-micaschists with large (0.5 to  $>1$  cm) garnets as well as amphibole-bearing rocks with large amphiboles ( $\geq 1$  cm). The SC rocks exposed belong to the marginal series (“bunte Randserie”) of intercalated garnet-micaschists, amphibolites, hornblende garbenschists, marbles and quartzites that delimit the main syncline of the SC. Further downhill, rocks of the central SC main syncline are encountered that are characterized by rather monotonous garnet-micaschists (“monotone Serie”). The SC–ÖC boundary is located a few meters uphill from a quartzite band that can unambiguously be assigned to the SC. The actual boundary



lies within several meters of gneisses containing small garnets and cannot unambiguously be localized in the field. A definite criterion for the distinction between SC and ÖC rocks, however, is the zoning pattern of garnets since SC rocks are monometamorphic and thus the garnets show continuous bell-shaped zoning patterns and the ÖC rocks are polymetamorphic thus the garnets show discontinuous chemical zoning.

#### 4.10 Field stop 10: Eclogites near Längenfeld, Variscan eclogite facies, Burgstein

The eclogites at the locality Burgstein belong to the central amphibolite complex in the ÖC and can be seen next to the tunnel portal. The central amphibolite complex is characterized by the occurrence of ultramafic rocks (peridotites), eclogites, amphibolites and rarely metacarbonates. The eclogites in this outcrop are Fe-eclogites, which appear dark-red and massive. The mineral assemblage is garnet (>50 vol%), omphacite, amphibole I, clinozoisite, quartz, apatite, pyrite and rutile. During subsequent alteration, symplectites (plagioclase + diopside) and amphibole II form (Miller & Thöny, 1995).

#### 4.11 Field stop 11: Köfels rock fall, Köfels

The Köfels rockslide is the biggest one in the Alps within basement rocks. It involved augengneisses of granitic composition as well as paragneisses. With a total mass of 3.2 km<sup>3</sup> of material (Brückl *et al.*, 2001) the rockslide was able to fill up the valley by some 100 meters. Due to the formation of a dam up to 100 m thick fluvio-lacustrine sediments were deposited in the Längenfelder basin (Heuberger, 1966, 1975). Relative, <sup>14</sup>C and




Fig. 6. Formation of the Tauern Window from the Early Miocene to the present (after Frisch *et al.*, 2000). Note that the sections are in E–W orientation.

cosmogenic isotope dating of exposed surfaces yielded an age of ~ 9800 cal BP for its activity (Ivy-Ochs *et al.*, 1998). After the main event at least one smaller event followed. The Köfels rockslide is well known for its frictionite, a pumice rock formed in the context of frictional melting (Erismann *et al.*, 1977).

## 5. References

- ACQUAFREDDA, P., BARGOSSO, G.M., CAGGIANELLI, A. & ROTTURA, A. (1997): Emplacement depths of Permian granitoids from central-eastern Southern Alps: estimates from hornblende-plagioclase thermobarometry. *Mineralogica et Petrographica Acta*, **40**: 45–53.
- ANGELMAIER, P., FRISCH, W. & DUNKL, I. (2000): Thermochronological constraints for the TRANSALP section of the Tauern Window. In *Abstr. 2<sup>nd</sup> International TRANSALP Colloquium*, Munich, 17.
- BARGOSSO, G. M., BONDI, M., LANDINI, F. & MORTEN, L. (1981): Il plutone di Monte Croce (Alto Adige, Nord Italia). *Rendiconti della Società Italiana di Mineralogia e Petrologia*, **38**: 155–162.
- BARGOSSO, G. M., ROTTURA, A., VERNIA, L., VISONÀ, D. & TRANNE, C. A. (1998): Guida all'escursione sul distretto vulcanico atesino e sulle plutoniti di Bressanone-Chiusa e Cima d'Asta. *Memorie della Società geologica italiana*, **53**: 23–41.
- BONIN, B., BRÄNDLEIN, P., BUSSY, F., DESMONS, J., EGGENBERGER, U., FINGER, F., GRAF, K., MARR, C., MERCOLLI, I., OBERHÄNSLI, R., PLOQUIN, A., VON QUADT, A., VON RAUMER, J.F., SCHALTEGGER, U., STEYRER, H.P., VISONÀ, D. & VIVIER, G. (1993): Late Variscan magmatic evolution of the Alpine basement. In von Raumer, J.F. & Neubauer, F. (eds): *The pre-Mesozoic geology in the Alps*. Springer, 169–199.
- BORSI, S., DEL MORO, A. & FERRARA, G. (1972): Età radiometriche delle rocce intrusive del Massiccio di Bressanone-Ivigna-Monte Croce (Alto Adige). *Bollettino della Società Geologica Italiana*, **91**: 387–406.
- BRANDNER, R., REITER, F. & TÖCHTERLE, A. (2008): Überblick zu den Ergebnissen der geologischen Vorerkundung für den Brenner-Basistunnel. *Geo.Alp*, **5**: 165–174.
- BRÜCKL, E., BRÜCKL, J. & HEUBERGER, H. (2001): Present structure and prefailure topography of the giant rockslide of Köfels. *Zeitschrift für Gletscherkunde und Glazialgeologie*, **37**: 49–79.



- CLIFF, R.A., DROOP, G.T.R. & REX, D.C. (1985): Alpine metamorphism in the south-east Tauern Window, Austria, II. Heating, cooling and uplift rates. *Journal of Metamorphic Geology*, **3**: 403–415.
- DACHS, E. (1986): High-pressure mineral assemblages and their breakdown products in metasediments South of the Grossvenediger, Tauern Window, Austria. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **66**: 145–161.
- DACHS, E. (1990): Geothermobarometry in metasediments of the southern Grossvenediger area (Tauern Window, Austria). *Journal of Metamorphic Geology*, **8**: 217–230.
- DAL PIAZ, G.V. & MARTIN, S. (1998): Evoluzione litosferica e magmatismo nel dominio austro-sudalpino dell'orogenesi varisca al rifting mesozoico. *Memorie della Società Geologica Italiana*, **53**: 43–62.
- DEL MORO, A. & VISONÀ, D. (1982): The epiplutonic Hercynian Complex of Bressanone (Brixen, Eastern Alps, Italy). *Neues Jahrbuch für Mineralogie, Abhandlungen*, **145**: 66–85.
- DINGELDEY, C., DALLMEYER, D.R., KOLLER, F. & MASSONNE, H.J. (1997): P-T-t history of the Lower Austroalpine Nappe Complex in the "Tarntaler Berge" NW of the Tauern Window: implications for the geotectonic evolution of the central Eastern Alps. *Contributions to Mineralogy and Petrology*, **129**: 1–19.
- DROOP, G.T.R. (1985): Alpine metamorphism in the south-east Tauern Window, Austria, I. P-T variations in space and time. *Journal of Metamorphic Geology*, **3**: 371–402.
- DROOP, G.T.R., LOMBARDO, B. & POGNATE, U. (1990): Formation and distribution of eclogite facies rocks in the Alps. In Carswell, D.A. (ed.): *Eclogite facies rocks*. Glasgow: Blackie, 225–259.
- ELIAS, J. (1998): The thermal history of the Ötztal-Stubai Complex (Tyrol, Austria, Italy) in the light of the lateral extrusion model. *Tübinger Geowissenschaftliche Arbeiten*, **42**: 172 p.
- ERISMANN, T., HEUBERGER, H. & PREUSS, E. (1977): Der Bimsstein von Köfels (Tirol), ein Bergsturz-"Friktionit". *Mineralogy and Petrology*, **24**: 67–119.
- EXNER, U., FUSSEIS, F., GRASEMANN, B., HABLER, G., LINNEN, M., SÖLVA, H., THIEDE, R. & THÖNI, M. (2001): Cretaceous eclogite-facies metamorphism in the Eastern Alps: New Insights, data and correlations from an interdisciplinary study. *Journal of Conference Abstracts*, **6**(1): 387.
- FRANK, W., MILLER, CH., PETRAKAKIS, K., PROCHASKA, K. & RICHTER, W. (1981): Das penninische Kristallin im Mittelabschnitt des Tauernfensters und die Rieserferner Intrusion mit ihrem Kontakthof. *Fortschritte der Mineralogie*, **59**: 97–128.
- FRANK, W., HÖCK, V. & MILLER, CH. (1987): Metamorphic and tectonic history of the central Tauern Window. In Flügel, H.W. & Faupl, R. (eds): *Geodynamics of the Eastern Alps*. Vienna: Deuticke, 34–54.
- FRANZ, G. & SPEAR, E.S. (1983): High pressure metamorphism of siliceous dolomites from the central Tauern Window, Austria. *American Journal of Science*, **283**: 396–413.
- FRISCH, W., DUNKL, I. & KUHLEMANN, J. (2000): Postcollisional orogenparallel large-scale extension in the Eastern Alps. *Tectonophysics*, **327**: 239–265.
- FÜGENSCHUH, B., SEWARD, D., MANCKTELOW, N. & FUMASOLI, M. (1997): Exhumation in a convergent orogen: the western Tauern window. *Terra Nova*, **9**: 213–217.
- FÜGENSCHUH, B., MANCKTELOW, N.S. & SEWARD, D. (2000): The Cretaceous to Neogene cooling and exhumation history of the Ötztal-Stubai basement complex, Eastern Alps: a structural and fission-track study. *Tectonics*, **19**: 905–918.
- GETTY, S.R. & SELVERSTONE, J. (1994): Stable isotopic and trace element evidence for restricted fluid migration in 2 GPa eclogites. *Journal of Metamorphic Geology*, **12**: 747–760.
- HABLER, G., LINNEN, M., THIEDE, R. & THÖNI, M. (2001): Eo-Alpine Andalusite in the Schneeberg Complex (Eastern Alps, Italy/Austria): Constraining the P-T-t-D path during Cretaceous metamorphism. *Journal of Conference Abstracts*, **6**: 340.
- HADITSCH, G. & MOSTLER, H. (1982): Zeitliche und stoffliche Gliederung der Erzvorkommen im Innsbrucker Quarzphyllit. *Geologisch-paläontologische Mitteilungen der Universität Innsbruck*, **12**: 1–40.
- HANDLER, R., GENSER, J., FRIEDL, G., HEIDORN, R., NEUBAUER, F., FRITZ, H. & TENCZER, V. (2000): Ar-Ar ages of white mica from Austroalpine basement units along the TRANSALP section: Tectonic implications. In *Abstr. 2nd International TRANSALP Colloquium*, Munich, 17.
- HERMANN, R.L., BLIKRA, L.H., NAUMANN, M., NILSEN, B., PANTHI, K.K., STROMEYER, D. & LONGVA, O. (2006): Examples of multiple rock slope collapses from Köfels (Ötztal valley, Austria) and western Norway. *Engineering Geology*, **83**: 94–108.
- HEUBERGER, H. (1966): Gletschergeschichtliche Untersuchungen in den Zentralalpen zwischen Sellrain und Ötztal. *Wissenschaftliche Alpenvereinshefte*, **20**: 1–126.
- HEUBERGER, H. (1975): Das Ötztal. Bergstürze und alte Gletscherstände, kulturgeographische Gliederung. *Innsbrucker Geographische Studien*, **2**: 213–249.
- HÖCK, V. & MILLER, CH. (1987): Mesozoic ophiolitic sequences and non-ophiolitic metabasites in the Hohe Tauern. In Flügel, H.W. & Faupl, E. (eds): *Geodynamics of the Eastern Alps*. Vienna: Deuticke, 16–33.
- HOINKES, G. (1973): Die Anatexis des Winnebachgranits (Ötztaler Alpen, Österreich) am Beispiel eines Aufschlusses. *Tschermaks Mineralogische und Petrographische Mitteilungen*, **20**: 225–239.
- HOINKES, G. & THÖNI, M. (1993): Evolution of the Ötztal-Stubai, Scarl-Campo and Ulten basement units. In von Raumer, J. F. & Neubauer, F. (eds): *Pre-Mesozoic geology in the Alps*. Heidelberg: Springer Verlag, 485–494.
- HOINKES, G., KOSTNER, A. & THÖNI, M. (1991): Petrologic constraints for Eo-Alpine eclogite facies metamorphism in the Austroalpine Ötztal Basement. *Mineralogy and Petrology*, **43**: 237–254.
- HOINKES, G., THÖNI, M., BERNHARD, F., KAINDL, R., LICHEM, C., SCHWEIGL, J., TROPPER, P. & COSCA, M. (1997): Metagranitoids and associated metasediments as indicators for the pre-Alpine magmatic and metamorphic evolution of the Western Austroalpine Ötztal Basement (Kauertal, Tirol). *Schweizerische Mineralogische und Petrographische Mitteilungen*, **77**: 299–314.
- HOINKES, G., KOLLER, F., RANTITSCH, G., DACHS, E., HÖCK, V., NEUBAUER, F. & SCHUSTER, R. (1999): Alpine metamorphism of the Eastern Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **79**: 155–181.
- HOLLAND, T.J.B. (1979): High water activities in the generation of high pressure kyanite eclogites in the Tauern Window, Austria. *Journal of Geology*, **87**: 1–27.



- HOLLAND, T.J.B. & RAY, N.J. (1985): Glaucofanite and pyroxene breakdown reactions in the Pennine units of the Eastern Alps. *Journal of Metamorphic Geology*, **3**: 417–438.
- HOLLAND, T.J.B. & RICHARDSON, S.W. (1979): Amphibole zonation in metabasites as a guide to the evolution of metamorphic conditions. *Contributions to Mineralogy and Petrology*, **70**: 143–148.
- HOSCHEK, G., KIRCHNER, E.C., MOSTLER, H. & SCHRAMM, J.M. (1980): Metamorphism in the Austroalpine units between Innsbruck and Salzburg (Austria) – a synopsis. *Mitteilungen der Österreichischen Geologischen Gesellschaft*, **71/72**: 335–341.
- IVY-OCHS, S., HEUBERGER, H., KUBIK, P. W., KERSCHNER, H., BONANI, G., FRANK, M. & SCHLÜCHTER, C. (1998): The age of the Köffels event. Relative,  $^{14}\text{C}$  and cosmogenic isotope dating of an early Holocene landslide in the Central Alps (Tyrol, Austria). *Zeitschrift für Gletscherkunde und Glazialgeologie*, **34**: 57–68.
- KLIER, R. (2005): Das Tarntal Mesozoikum: Petrologie und Geologie einer enigmatischen Einheit in den Ostalpen. Unpubl. MSc Thesis, Univ. of Innsbruck, Austria, 99 p.
- KLÖTZLI-CHOWANETZ, E., KLÖTZLI, U. & KOLLER, F. (1997): Lower Ordovician migmatization in the Ötztal crystalline basement (Eastern Alps, Austria): linking U-Pb and Pb-Pb dating with zircon morphology. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **77**: 315–324.
- KOLENPRAT, B. (1998): Bericht 1997 über geologische Aufnahmen im Altkristallin und Unterostalpin im Raum Glungezer auf Blatt 148 Brenner. *Jahrbuch der Geologischen Bundesanstalt*, **141**: 292.
- KOLENPRAT, B., ROCKENSCHAUB, M. & FRANK, W. (1999): The tectonometamorphic evolution of Austroalpine units in the Brenner Area (Tirol, Austria) - structural and tectonic implications. *Tübinger Geowissenschaftliche Arbeiten, Ser. A*, **52**: 116–117.
- KONZETT, J. & HOINKES, G. (1991): High pressure metamorphism in the Austroalpine Schneeberg Complex and adjacent Ötztal basement rocks. *Terra Abstracts*, **3**: 88.
- KONZETT, J. & HOINKES, G. (1996): Paragonite-hornblende assemblages and petrological significance: an example from the Austroalpine Schneeberg Complex, Southern Tyrol, Italy. *Journal of Metamorphic Geology*, **14**: 85–101.
- KRALIK, M., KRUMM, H. & SCHRAMM, J.M. (1987): Low grade and very low grade metamorphism in the Northern Calcareous Alps and in the Greywacke Zone: Illite crystallinity data and isotopic ages. In Flügel, H.W. & Faupl, P. (eds): *Geodynamics of the Eastern Alps*. Vienna: Deuticke, 164–178.
- MILLER, CH. (1974): On the metamorphism of the eclogites and high-grade blueschists from the Penninic Terrane of the Tauern Window, Austria. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **54**: 371–384.
- MILLER, CH. (1977): Chemismus und phasenpetrologische Untersuchungen der Gesteine aus der Eklogitzone des Tauernfensters, Österreich. *Tschermaks Mineralogische und Petrographische Mitteilungen*, **24**: 221–277.
- MILLER, CH. (1986): Alpine high-pressure metamorphism in the Eastern Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **66**: 139–144.
- MILLER, CH. (1987): Die Eklogite des Tauernfensters. *Field Guide, Tauerntagung, Mittersill*, 10 p.
- MILLER, CH. & THÖNI, M. (1995): Origin of eclogites from the Austroalpine Ötztal basement (Tirol, Austria): geochemistry and Sm-Nd vs. Rb-Sr isotope systematics. *Chemical Geology*, **122**: 199–225.
- MOSTLER, H. (1973): Alter und Genese ostalpinen Spatmagnetit unter besonderer Berücksichtigung der Magnetitlagerstätten im Westabschnitt der Nördlichen Grauwackenzone (Tirol, Salzburg). *Veröffentlichungen der Universität Innsbruck*, **86**: 237–266.
- NEUBAUER, F., HOINKES, G., SASSI, F. P., HANDLER, R., HÖCK, V., KOLLER, F. & FRANK, W. (1999): Pre-Alpine metamorphism in the Eastern Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **79**: 41–62.
- OBERHÄNSLI, R., BOUSQUET, R., ENGI, M., GOFFÉ, B., GOSSO, G., HANDY, M., KOLLER, F., LARDEAUX, J.-M., POLINO, R., ROSSI, P., SCHUSTER, R., SCHWARTZ, S., SPALLA, I. E., AGARD, P., BABIST, J., BERGER, A., BERTLE, R., BUCHER, K., BURRI, T., HEITZMANN, P., HOINKES, G., JOLIVET, L., KELLER, L., LINNEN, M., LOMBARDO, B., MARTINOTTI, G., MICHARD, A., PESTAL, G., PROYER, A., RANTISCH, G., ROSENBERG, C., SCHRAMM, J., SÖLVA, H., THÖNI, M. & ZUCALI, M. (2004): Map of the metamorphic structure of the Alps 1:1 000 000. Commission for the Geological Map of the World: Paris.
- PIBER, A. (2005): Tectonometamorphic evolution of the Austro-Alpine nappes in the northern Zillertal-Area, Eastern Alps. Unpubl. PhD Thesis, Univ. of Innsbruck, Austria, 268 p.
- PIBER, A. & TROPPER, P. (2005): The polymetamorphic evolution of the Austroalpine Innsbruck Quartzphyllite Complex. *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, 151.
- PIBER, A. & TROPPER, P. (2007): Muscovite-chlorite-quartz thermobarometry of the Permian metamorphic overprint of the central part of the western Innsbruck Quartzphyllite Complex (Tyrol, Austria). *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, **153**: 291–300.
- PIBER, A., TROPPER, P. & MIRWALD, P.W. (2008): The metamorphic evolution of the Patscherkofel Crystalline Complex (Tyrol, Eastern Alps, Austria). *Austrian Journal of Earth Sciences*, **101**: 27–35.
- POMELLA, H., FÜGENSCHUH, B. & STIPP, M. (2008): The Giudicarie Fault System at the transition between Southern and Eastern Alps (Northern Italy) A new structural analysis. *Abstr. Vol., 6th Swiss Geoscience Meeting, Lugano 2008*, 55.
- PURTSCHALLER, F. (1969): Petrographische Untersuchungen an Aluminosilikatgneisen des Ötztals-Stubaier Altkristallins. *Tschermaks Mineralogische und Petrographische Mitteilungen*, **13**: 35–54.
- RAITH, M., MEHRENS, CH. & THÖLE, W. (1980): Gliederung, tektonischer Bau und metamorphe Entwicklung der penninischen Serien im südlichen Venediger-Gebiet, Osttirol. *Jahrbuch der Geologischen Bundesanstalt*, **123**: 1–37.
- RING, U. & RICHTER, C. (1994): The Variscan structural and metamorphic evolution of the eastern Southalpine basement. *Journal of Geological Society, London*, **151**: 755–766.
- ROCKENSCHAUB, M. (1998): Bericht 1997 über geologische Aufnahmen in der Quarzphyllitzone und im Tauernfenster auf Blatt 148 Brenner. *Jahrbuch der Geologischen Bundesanstalt*, **141**: 295.



- ROCKENSCHAUB, M., KOLENPRAT, B. & FRANK, W. (1999): The tectonometamorphic evolution of Austroalpine units in the Brenner area (Tirol, Austria) - new geochronological implications. *Tübinger Geowissenschaftliche Arbeiten*, **52**: 118–119.
- ROTH, R. (1983): Petrographie und Tektonik der mittelostalpinen Kellerjochgneis-Decke und angrenzender Gebiete zwischen Schwaz und Märzengrund (Tirol.). Unpubl. PhD Thesis, Univ. of Münster, Germany, 196 p.
- ROTTURA, A., BARGOSSO, G.M., CAGGIANELLI, A., DEL MORO, A., VISONÀ, D. & TRANNE, C.A. (1998): Origin and significance of the Permian high-K alkaline magmatism in the central-eastern Southern Alps, Italy. *Lithos*, **45**: 329–348.
- SASSI, F. P. & SPIESS, R. (1992): Further data on pre-Alpine metamorphic pressure conditions of the Austridic phyllitic complexes in the Eastern Alps. In Carmignagi, L. & Sassi, F.P. (eds): Contributions to the geology of Italy with special regard to the Paleozoic basements. IGCP Project 276, Newsletter **5**: 297–307.
- SASSI, F.P. & SPIESS, R. (1993): The South Alpine metamorphic basement in the Eastern Alps. In von Raumer, J.F. & Neubauer, F. (eds): Pre-Mesozoic geology in the Alps. Heidelberg: Springer, 599–607.
- SCHMID, S.M., FÜGENSCHUH, B., KISSLING, E. & SCHUSTER, R. (2004): Tectonic map and overall architecture of the Alpine orogen, *Eclogae Geologicae Helvetiae*, **97/1**: 93–117.
- SCHMIDEGG, O. (1964): Die Ötztaler Schubmasse und ihre Umgebung. *Verhandlungen der Geologischen Bundesanstalt*, 1964(1), 24–47.
- SCHMIDLECHNER, M., NEUBAUER, F. & HANDLER, R. (2006): Extent and age of metamorphism of the central Grauwacken Zone, Eastern Alps: An Ar-Ar study. *Pangeo 2006*, Innsbr. Univ. Press, 314–315.
- SCHUSTER, R., SCHARBERT, S., ABART, R. & FRANK, W. (2001): Permian-Triassic extension and related HAT/LP metamorphism in the Austroalpine – Southalpine realm. *Mitteilungen der Gesellschaft der Geologie- und Bergbaustudenten in Österreich*, **45**: 112–141.
- SCHUSTER, R., KOLLER, F., HÖCK, V., HOINKES, G. & BOUSQUET, R. (2004): Explanatory notes to the map: metamorphic structure of the Alps, metamorphic evolution of the Eastern Alps. *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, **149**: 175–199.
- SCHWEIGL, J. (1993): Kristallingeologische Untersuchungen in den Nauderer Bergen (Westliche Ötztaler Alpen, Tirol). Unpubl. MSc Thesis, University of Vienna, Austria, 87 p.
- SCHWEIGL, J. (1995): Neue geochronologische und isotopengeologische Daten zur voralpidischen Entwicklungsgeschichte im Ötztal-kristallin (Ostalpen). *Jahrbuch der Geologischen Bundesanstalt*, **138**(1): 131–149.
- SELVERSTONE, J. (1993): Micro- to macroscale interactions between deformational and metamorphic processes, Tauern Window, Eastern Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen* **73**: 229–239.
- SELVERSTONE, J., FRANZ, G., THOMAS, S. & GETTY, S. (1992): Fluid variability in 2GPa eclogites as an indicator of fluid behaviour during subduction. *Contributions to Mineralogy and Petrology* **112**: 341–357.
- SÖLLNER, F. (1982): Zur Altersstellung des Winnebach-Migmatits im Ötztal (Ostalpen). *Verhandlungen der Geologischen Bundesanstalt*, **2**: 95–106.
- SÖLLNER, F. & SCHMIDT, K. (1981): Rb/Sr- und U/Pb-Datierungen am Winnebach-Migmatit (Ötztaler Alpen, Österreich). *Fortschritte der Mineralogie*, **59**, Beiheft 1: 188.
- SÖLVA, H., THÖNI, M., GRASEMAN, B. & LINNEN, M. (2001): Emplacement of eo-Alpine high-pressure rocks in the Austroalpine Ötztal complex (Texel group, Italy/Austria). *Geodinamica Acta*, **14**: 345–360.
- SÖLVA, H., GRASEMANN, B., THÖNI, M., THIEDE, R. & HABLER, G. (2005): The Schneeberg Normal Fault Zone: normal faulting associated with Cretaceous SE-directed extrusion in the Eastern Alps, *Tectonophysics*, **401**: 143–166.
- SPEAR, E.S. & FRANZ, G. (1986): P-T evolution of metasediments from the Eclogite Zone, south-central Tauern Window, Austria. *Lithos*, **19**: 219–234.
- SPIESS, R. (1995): The Passeier-Jaufen Line: a tectonic boundary between Variscan and eo-Alpine Meran-Mauls basement. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **75**: 413–425.
- THÖNI, M. (1981): Degree and evolution of the Alpine metamorphism in the Austroalpine unit west of the Hohe Tauern in the light of K/Ar and Rb/Sr age determinations on micas. *Jahrbuch der Geologischen Bundesanstalt*, **124**: 111–174.
- THÖNI, M. (1986): The Rb-Sr thin slab isochron method – an unreliable geochronologic method for dating geologic events in polymetamorphic terrains? *Memorie di Scienze Geologiche*, **38**: 283–352.
- THÖNI, M. (1999): A review of geochronological data from the Eastern Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **79**: 209–230.
- THÖNY, W.F. (2008): Permian contact metamorphism in the Southalpine: U-Pb and U-Th-Pb geochronology of plutons and contact metamorphism. Unpubl. PhD Thesis, Univ. of Innsbruck, Austria, 200 p.
- THÖNY, W.F., TROPPER, P., SCHENNACH, F., KRENN, E., FINGER, F., KAINDL, R., BERNHARD, F. & HOINKES, G. (2008): The metamorphic evolution of migmatites from the Ötztal Complex (Tyrol, Austria) and constraints on the timing of the pre-Variscan high-T event in the Eastern Alps. *Swiss Journal of Geosciences*, **101**: 111–126.
- TROPPER, P. & HOINKES, G. (1996): Thermobarometry in Al<sub>2</sub>SiO<sub>5</sub>-bearing metapelites in the western Austroalpine Ötztal-basement. *Mineralogy and Petrology*, **58**: 145–170.
- VIOLA, G. & O'DONNELL, F., MANCKTELOW, N.S. (2004): Analogue modelling of reverse fault reactivation in strike-slip and transpressive regimes: application to the Giudicarie fault system, Italian Eastern Alps. *Journal of Structural Geology*, **36**: 401–418.
- ZIMMERMANN, R., HAMMERSCHMIDT, K. & FRANZ, G. (1994): Eocene high pressure metamorphism in the Penninic units of the Tauern Window (Eastern Alps). Evidence from <sup>40</sup>Ar-<sup>39</sup>Ar dating and petrological investigations. *Contributions to Mineralogy and Petrology*, **117**: 175–186.
- ZWINGMANN, H. & MANCKTELOW, N.S. (2004): Timing of Alpine fault gouges. *Earth and Planetary Science Letters*, **223**: 415–423.



## Appendix – Itinerary for IMA2010 AT2 Field trip

### Thursday, August 19, 2010 (Day 1). Transect through the Eastern Alps (deformational focus)

06.00–13.00	Travel to Innsbruck and lunch
13.00–14.00	Field stop 1: Sillschlucht: introduction to geological background
14.00–16.00	Field stops 2, 3 and 4: Stefansbrücke, Nösslach, Brenner, Brenner fault zone
16.00–17.00	Field stops 5 and 6: Franzensfeste, Mauls, Permian and Oligocene intrusions Accommodation in Sterzing

### Friday, August 20, 2010 (Day 2). Transect through the Eastern Alps (metamorphic focus)

09.00–10.00	Field stop 7: Jaufen pass, Cretaceous tectonics
10.00–11.00	Field stop 8: Saltaus, Cretaceous eclogite-facies metamorphism
11.00–12.00	Field stop 9: Timmelsjoch: Ötztal Complex, Schneeberg Complex
12.00–13.00	Lunch break at the Timmelsjoch
13.00–14.00	Field stop 10: Burgstein: Variscan eclogites
14.00–15.00	Field stop 11: Köfels, rock fall
15.00–22.00	Travel to Budapest



X 175 526







## INSTRUCTIONS FOR AUTHORS

### GENERAL

Acta Mineralogica-Petrographica (AMP) publishes articles (papers longer than 4 printed pages but shorter than 16 pages, including figures and tables), notes (not longer than 4 pages, including figures and tables), and short communications (book reviews, short scientific notices, current research projects, comments on formerly published papers, and necrologies of 1 printed page) dealing with crystallography, mineralogy, ore deposits, petrology, volcanology, geochemistry and other applied topics related to the environment and archaeometry. Articles longer than the given extent can be published only with the prior agreement of the editorial board.

In the form of two subseries, AMP publishes materials of conferences (AMP Abstract Series) and field guides (AMP Field Guide Series), or, occasionally supplement issues related to other scientific events.

The journal accepts papers that represent new and original scientific results, which have not appeared elsewhere before, and are not in press either.

All articles and notes submitted to AMP are reviewed by two referees (short communications will be reviewed only by one referee) and are normally published in the order of acceptance, however, higher priority may be given to Hungarian researches and results coming from the Alpine-Carpathian-Dinaric region. Of course, the editorial board does accept papers dealing with other regions as well, let they be compiled either by Hungarian or foreign authors.

The manuscripts (prepared in harmony of the instructions below) must be submitted to the Editorial Office in triplicate. All pages must carry the author's name, and must be numbered. At this stage (revision), original illustrations and photographs are not required, though, quality copies are needed. It is favourable, if printable manuscripts are sent on disk, as well. In these cases the use of Microsoft Word or any other IBM compatible editing programmes is suggested.

### LANGUAGE

The language of AMP is English.

### PREPARATION OF THE MANUSCRIPT

The different parts of the manuscript need to meet the instructions below:

#### Title

The title has to be short and informative. No subtitles if possible. If the main title is too long, an additional shortened title is needed for the running head.

#### Author

The front page has to carry (under the main title) the full name(s) (forename, surname), affiliation(s), current address(es), e-mail address(es) of the author(s).

#### Abstract and keywords

The abstract is required to be brief (max. 250 words), and has to highlight the aims and the results of the article. The abstracts of notes are alike (but max. 120 words). As far as possible, citations have to be avoided.

The abstract has to be followed by 4 to 10 keywords.

#### Text and citations

The format of the manuscripts is required to be: double-spacing (same for the abstract), text only on one side of the page, size 12 Times New Roman fonts. Margin width is 2.5 cm, except the left margin, which has to be 3.5 cm wide. Underlines and highlights ought not be used. Please avoid the use of foot and end notes. Accents of Romanian, Slovakian, Czech, Croatian etc. characters must be marked on the manuscript clearly.

When compiling the paper an Introduction – Geological setting – Materials and Methods – Results – Conclusions structure is suggested.

The form of citations is: the author's surname followed by the date of publication e.g. (Szederkényi, 1996). In case of two authors: (Rosso and Bodnar, 1995) If there are more than two authors, after the first name the co-authors must be denoted as "et al.", e.g. (Roser et al., 1980).

### REFERENCES

The reference list can only consist of published papers, M.Sc., Ph.D. and D.Sc. theses, and papers in press.

Only works cited previously in the text can be put in the reference list. Examples:

Best, M.G., Christiansen, E.H. (eds.) (2001): Igneous petrology. Blackwell, Edinburgh, 458p.

Upton, B.G.J., Emeleus, C.H. (1987): Mid-Proterozoic alkaline magmatism in southern Greenland: the Gardar province. In: Fitton, J.G., Upton, B.G.J. (eds.): Alkaline Igneous Rocks. Blackwell, Edinburgh, 449–472.

Rosso, K.M., Bodnar, R.J. (1995): Microthermometric and Raman spectroscopic detection limits of CO<sub>2</sub> in fluid inclusions and the Raman spectroscopic characterization of CO<sub>2</sub>. *Geochimica et Cosmochimica Acta*, **59**, 3961–3975.

Szederkényi, T. (1996): Metamorphic formations and their correlation in the Hungarian part of Tisia Megaunit (Tisia Megaunit Terrane). *Acta Mineralogica-Petrographica*, **37**, 143–160.

Bakker, R.J. (2002): <http://www.unileoben.ac.at/~buero62/minpet/Ronald/Programs/Computer.html>. Accessed: June 15, 2003.

The full titles of journals ought to be given. In case more works of the same author are published in the same year, then these has to be differentiated by using a, b, etc. after the date.

### ILLUSTRATIONS

Finally, each figure, map, photograph, drawing, table has to be attached in three copies, they must be numbered and carry the name of the author on their reverse. All the illustrations ought to be printed on separate sheets, captions as well if possible. Foldout tables and maps are not accepted. In case an illustration is not presented in digital form then one of the copies has to be submitted as glossy photographic print suitable for direct reproduction. Photographs must be clear and sharp. The other two copies of the illustrations can be quality reproductions. Coloured figure, map or photograph can only be published at the expense of the author(s).

The width of the illustrations can be 56, 87, 118, or 180 mm. The maximum height is 240 mm (with caption).

All figures, maps, photographs and tables are placed in the text, hence, it is favourable if in case of whole page illustrations enough space is left on the bottom for inserting captions. In the final form the size of the fonts on the illustrations must be at least 1,5 mm, their outline must be 0,1 mm wide. Digital documents should be submitted in JPG-format. The resolution of line-drawings must be 400 dpi, while that of photographs must be 600 dpi. The use of Corel Draw for preparing figures is highly appreciated, and in this case please submit the .CDR file, as well.

### PROOFS AND OFFPRINTS

After revision the author(s) receive only the page-proof. The accepted and revised manuscripts need to be returned to the Editors either on disc, CD or as an e-mail attachment. Proofreading must be limited to the correction of typographical errors. If an illustration cannot be presented in digital form, it must be submitted as a high quality camera-ready print.

The author(s) will receive 25 free offprints. On payment of the full price, further offprints can be ordered when the corrected proofs are sent back.

#### Manuscripts for publication in the AMP should be submitted to:

Dr. Pál-Molnár Elemér

E-mail: [palm@geo.u-szeged.hu](mailto:palm@geo.u-szeged.hu)

Dr. Batki Anikó

E-mail: [batki@geo.u-szeged.hu](mailto:batki@geo.u-szeged.hu)

Phone: 00-36-62-544-683, Fax: 00-36-62-426-479

Department of Mineralogy, Geochemistry and Petrology University of Szeged  
P.O. Box 651

H-6701 Szeged, Hungary

Distributed by the Department of Mineralogy, Geochemistry and Petrology,  
University of Szeged, Szeged, Hungary.





# ACTA MINERALOGICA-PETROGRAPHICA FIELD GUIDE SERIES

VOLUME 2 2010

HU ISSN 0324-6523

HU ISSN 2061-9766

## MAP OF THE IMA2010 FIELD TRIP AT2



Map provided by the Stockwork Exploration Ltd. ([www.stockwork.eu](http://www.stockwork.eu))